



Theorems		For Your FOLDABLE
Special Segments of Similar Triangles		
<p>7.8 If two triangles are similar, the lengths of corresponding altitudes are proportional to the lengths of corresponding sides.</p> <p>Abbreviation $\sim\Delta s$ have corr. altitudes proportional to corr. sides.</p> <p>Example If $\triangle ABC \sim \triangle FGH$, then $\frac{AD}{FJ} = \frac{AB}{FG}$.</p>		
<p>7.9 If two triangles are similar, the lengths of corresponding angle bisectors are proportional to the lengths of corresponding sides.</p> <p>Abbreviation $\sim\Delta s$ have corr. \angle bisectors proportional to corr. sides.</p> <p>Example If $\triangle KLM \sim \triangle QRS$, then $\frac{LP}{RT} = \frac{LM}{RS}$.</p>		
<p>7.10 If two triangles are similar, the lengths of corresponding medians are proportional to the lengths of corresponding sides.</p> <p>Abbreviation $\sim\Delta s$ have corr. medians proportional to corr. sides.</p> <p>Example If $\triangle ABC \sim \triangle WXY$, then $\frac{CD}{YZ} = \frac{AB}{WX}$.</p>		


1. In the figure $\triangle ABC \sim \triangle FDG$. Find the value of x .



2. The triangles below are similar. Find the value of x .



3. The triangles below are similar. Find the value of x .



Theorem 7.11 **For Your FOLDABLE**


Triangle Angle Bisector

An angle bisector in a triangle separates the opposite side into two segments that are proportional to the lengths of the other two sides.


Example If \overline{JM} is an angle bisector of $\triangle JKL$,
 then $\frac{KM}{LM} = \frac{KJ}{LJ}$.
← segments with vertex K
← segments with vertex L

4. Find x in the following diagrams:


a.



c.



b.



d.

